

AperTO - Archivio Istituzionale Open Access dell'Università di Torino

Fossil amphibians and reptiles from Plakias, Crete: a glimpse into the earliest late Miocene herpetofaunas of southeastern Europe

This is the author's manuscript

Original Citation:

Availability:

This version is available <http://hdl.handle.net/2318/1606927> since 2017-05-19T15:16:14Z

Published version:

DOI:<http://dx.doi.org/doi:10.1016/j.geobios.2016.09.004>

Terms of use:

Open Access

Anyone can freely access the full text of works made available as "Open Access". Works made available under a Creative Commons license can be used according to the terms and conditions of said license. Use of all other works requires consent of the right holder (author or publisher) if not exempted from copyright protection by the applicable law.

(Article begins on next page)

This Accepted Author Manuscript (AAM) is copyrighted and published by Elsevier. It is posted here by agreement between Elsevier and the University of Turin. Changes resulting from the publishing process - such as editing, corrections, structural formatting, and other quality control mechanisms - may not be reflected in this version of the text. The definitive version of the text was subsequently published in GEOBIOS, None, 9999, <http://dx.doi.org/doi:10.1016/j.geobios.2016.09....>

You may download, copy and otherwise use the AAM for non-commercial purposes provided that your license is limited by the following restrictions:

- (1) You may use this AAM for non-commercial purposes only under the terms of the CC-BY-NC-ND license.
- (2) The integrity of the work and identification of the author, copyright owner, and publisher must be preserved in any copy.
- (3) You must attribute this AAM in the following format: Creative Commons BY-NC-ND license (<http://creativecommons.org/licenses/by-nc-nd/4.0/deed.en>), <http://dx.doi.org/doi:10.1016/j.geobios.2016.09...>

When citing, please refer to the published version.

Link to this full text:

<http://hdl.handle.net/2318/1606927>

**Fossil amphibians and reptiles from Plakias, Crete: a glimpse into the
earliest late Miocene herpetofaunas of southeastern Europe**

Georgios L. Georgalis^{1,2,*}, Andrea Villa², Evangelos Vlachos^{3,4}, Massimo Delfino^{2,5}

¹ Department of Geosciences, University of Fribourg / Freiburg, Chemin du Musée 6, 1700
Fribourg, Switzerland

² Dipartimento di Scienze della Terra, Università di Torino, Via Valperga Caluso 35, 10125
Torino, Italy

³ CONICET and Museo Paleontologico Egidio Feruglio, Av. Fontana 140, 9100 Trelew,
Chubut, Argentina

⁴ School of Geology, Aristotle University of Thessaloniki, University Campus, 54124
Thessaloniki, Greece

⁵ Institut Català de Paleontologia Miquel Crusafont, Universitat Autònoma de Barcelona,
(UAB), Edifici Z, Carrer de les Columnes s/n, Campus de la UAB, 08193 Cerdanyola del
Vallès, Barcelona, Spain.

* corresponding author. E-mail address: dimetrodon82@gmail.com (G.L. Georgalis)

Abstract

Fossil amphibians and reptiles from the earliest late Miocene (early Tortonian, MN 9) of Plakias, Crete, Greece, are described in this paper. Most of the material is fragmentary, precluding precise taxonomic assignment. Nevertheless, the herpetofauna of Plakias is here shown to be diverse, comprising at least six different taxa: one alytid anuran, one crocodylian, two turtles (one pan-trionychid and one geoemydid) and two squamates (one amphisbaenian and one colubroid snake). The crocodylian material represents the first such fossils described from Greece and furthermore, one of the latest occurrences of this group in Europe. The pan-trionychid and the geoemydid represent the oldest occurrences of these groups in Greece and further add to their scarce Miocene record from the country. The first description of a fossil amphisbaenian from Greece is also provided. The new specimens from Plakias add to our knowledge of the Miocene herpetofaunas of southeastern Europe.

Keywords:

Anura

Testudines

Crocodylia

Amphisbaenia

Serpentes

Miocene

1. Introduction

Miocene herpetofaunas from Europe are relatively well understood (Rage and Augé, 1993; Sanchiz, 1998; Rage and Roček, 2003; Szyndlar and Rage, 2003; Augé, 2005; Danilov, 2005). However, the focus of previous studies is heavily unbalanced towards the herpetofaunas of the western and central parts of the continent, whereas only few papers have dealt with the eastern European ones (Szyndlar and Zerova, 1990; Szyndlar, 1991a, b; Rage and Augé, 1993; Antunes, 1994; Rage and Roček, 2003; Danilov, 2005; Daza et al., 2014). On the other hand, even less is known about the Miocene amphibians and reptiles from southeastern Europe, despite the fact that this region played a pivotal biogeographic role during that period: several dispersals events from both Africa and Asia have taken place during this time interval (Rögl, 1999; Koufos et al., 2005). As such, the study of southeastern European localities could clarify significant aspects about the biogeography, evolutionary history and extinction events of certain European groups.

Greece harbors a large number of Miocene localities, of which several have been well known and studied for more than a century (Koufos, 2006, and references therein). However, mammal finds were almost always the main focus of these studies, thereby neglecting other important tetrapod groups, such as amphibians and reptiles. Nevertheless, important fossil finds during the last 160 years have shown a diverse array of Miocene reptiles (Gaudry, 1862-1867; Römer, 1870; Weithofer, 1888; Szalai, 1931; Paraskevaidis, 1955; Bachmayer, 1967; Richter, 1995; Szyndlar, 1995; Georgalis et al., 2013; Georgalis and Kear, 2013; Vlachos and Tsoukala, 2014; Vlachos et al., 2015b; Garcia et al., 2016; Georgalis et al., 2016b), whereas amphibian remains are practically unknown from this time interval from Greece, with only few, sporadic, published occurrences (Sanchiz, 1998; Rage and Roček, 2003).

Here we describe new amphibian and reptile finds from the earliest late Miocene (early Tortonian, MN 9) locality of Plakias, on the Island of Crete, southern Greece. The

fragmentary nature of these fossils precludes any taxonomic designation to the species level. Higher level taxonomic determination of the specimens was possible, however, revealing an unexpectedly diverse assemblage consisting of anurans, pan-trionychids, geoemydids, crocodylians, amphisbaenians, and snakes.

Institutional Abbreviations: NHMC, Natural History Museum and University of Crete, Greece; UU, Department of Earth Sciences, University of Utrecht, The Netherlands.

2. Geological framework

All specimens described herein were collected from the locality of Plakias (also known under the spelling Plakia). This is an earliest late Miocene locality, situated in the Agios Vasileios municipality, Rethymnon periphery, Island of Crete (Fig. 1). Its exact age was initially believed to be younger than late Astaracian (late MN 7 / MN 8) (de Bruijn and Meulenkamp, 1972), a suggestion that was tentatively followed by van der Made (1996) and Koufos (2006). However, it was recently shown that the fauna pertains to the Vallesian (early Tortonian, MN 9), with an age of approximately 9.9 Ma (de Bruijn et al., 2012; Koufos and Kostopoulos, 2013). The Plakias Basin is filled with continental deposits characterized by alternations of silty clays, silts, sandstones and conglomerates (de Bruijn et al., 2012). Fossils originate from grayish clays overlying beige to brown clays with calcretes in a shallow gully complex, southeast of Plakias (de Bruijn et al., 2012).

The palaeoherpetofauna of Plakias cannot confirm, neither dispute with certainty the suggested age of the locality as Vallesian (MN 9), earliest late Miocene (de Bruijn et al., 2012). However, it is noted that pan-trionychids and crocodylians are better represented in early and middle Miocene localities of Europe, rather than late Miocene ones (Karl, 1999; Delfino and Rossi, 2013). However, since none among the Plakias representatives of the

herpetofauna can be attributed to the species level, no further age correlations can be safely made. In any case, the age information on the basis of micromammals allows referring the whole faunal assemblage to the earliest late Miocene (MN 9) (de Bruijn et al., 2012).

3. Material and methods

All the amphibians, crocodylians and squamates, and part of the pan-trionychid and geoemydid material described in this study is housed in the collections of the University of Utrecht (UU). This amphibian and reptile material was collected along with the fossil micromammals that were described by Bruijn and Meulenkamp (1972) from the first period of field work at Plakias.

Part of the geoemydid and the pan-trionychid material belongs to the collections of the Natural History Museum of Crete (NHMC) and represents material that was collected by S. Kuss in the 1970's. This material was originally in the Geologisch-Paläontologisches Institut der Universität Freiburg, Germany, and later formed part of the collections of the Natural History Museum of Karlsruhe, Germany, before its recent return to the NHMC. This material was studied in the doctoral thesis of one of us (Vlachos, 2015) and is further presented here in the context of the whole herpetofauna of Plakias. It is not possible to determine the exact stratigraphic origin of the material collected by Kuss, as he did not publish anything about it. It has been recently noted, however, that the old (de Bruijn and Meulenkamp, 1972) and new (de Bruijn et al., 2012) micromammal collections of Plakias could “come from slightly different stratigraphic levels” (de Bruijn et al., 2012: p. 61) due to building activity in the area during the last 40 years. However, the same authors noted that “the 25 cm thick grayish silty clay sampled in 2010 seems to be the only bed that contains vertebrate remains” (de Bruijn et al., 2012: p. 61). Following this line of reasoning, and in the absence of conflicting evidence, we treat all the material described herein as originating from the same site.

Taxonomy follows Pyron and Wiens (2011) for anurans, Joyce et al. (2004) for turtles, Brochu (2000) for crocodylians, and Gauthier et al. (2012), Pyron et al. (2014), and Wallach et al. (2014) for squamates. Anatomical terminology follows Sanchiz (1998) for anurans, Zangerl (1969) for turtles, Steel (1973) for crocodylians, Estes (1983) for amphisbaenians, and Rage (1984) for snakes.

4. Systematic Palaeontology

AMPHIBIA Linnaeus, 1758

ANURA Fischer von Waldheim, 1813

ALYTIDAE Fitzinger, 1843

cf. Alytidae indet.

Referred specimens: UU PL 701, a fragmentary trunk vertebra; UU PL 702, a single, partial tibiofibula.

Description: UU PL 701 (Fig. 2): This trunk vertebra preserves only the centrum and, on both sides, a small, basal portion of the neural arch. The centrum is roughly 2.8 mm long and slightly dorsoventrally flattened. The condyle bears a small, but well-defined condylar neck that is typical of opisthocoelous vertebrae and therefore characterizes the anterior extremity of the centrum. The anterior condyle and the posterior cotyle are rather subcircular.

UU PL 702: The preservational status of this tibiofibula fragment is poor. It preserves only a terminal portion, showing the presence of the two fused elements. Moreover, the fact that this skeletal element has limited diagnostic value hinders a precise identification.

Remarks: UU PL 701 can be tentatively referred to the Alytidae on the basis of the condylar neck that marks the condyle. This character has not been described in the literature (e.g. Bailon, 1999) but in our experience, this trait is characteristic for this group of frogs, which at least in some cases have also a comparable size. UU PL 702 is here tentatively referred to the

146 same taxon represented by the co-occurring vertebra. Most probably, the amphibian material
147 presented herein is the same that Sanchiz (1998: p. 168) referred to it as “Platkia (sic), Greece,
148 Neogene: Discoglossinae indet. (Sanchiz, unpublished)” (Borja Sanchiz, pers. commun. to
149 GLG, February 2016).

150 It is beyond the scope of this paper to evaluate the taxonomic affinities of European
151 painted frogs and we follow Frost et al. (2006) in using the term Alytidae for all these
152 amphibians. Despite their confined extant distribution, alytids were once widespread in
153 Europe (Rage and Roček, 2003). In Greece, other fossils attributed to this group have been
154 briefly reported from the Miocene of Aliveri, Ano Metochi, Biodrak, Lefkon, Maramena,
155 Monasteri, and Pikermi, the Pliocene of Kardina, Ptolemais, and Spilia and the Miocene or
156 Pliocene of Rema Aslan and Rema Marmara (Rage and Roček, 2003).

157

158 REPTILIA Laurenti, 1768

159 TESTUDINES Batsch, 1788

160 PAN-TRIONYCHIDAE Joyce et al., 2004

161 Pan-Trionychidae indet.

162 **Referred specimens:** UU PL 703, costal fragment; UU PL 704, costal fragment; NHMC
163 21.7.3.1670, neural; NHMC 21.7.3.1671, plastron fragment.

164 **Description:** NHMC 21.7.3.1670 (Fig. 3A): This specimen corresponds to an almost
165 complete neural. Although the specimen is eroded, the shape of the neural appears to be
166 hexagonal with short lateral sides. Dorsally, the distinctive sculpturing is visible, consisting
167 mainly of small pits. Viscerally, the attachment for the vertebra is preserved.

168 UU PL 703 (Fig. 3B): This specimen corresponds to a fragment of a costal, as is shown by the
169 presence of a rib on the visceral part. On the distal side, a part of the rib is apparent as well.
170 Dorsally, the distinctive sculpturing is visible, consisting of small pits that are mainly

separated from one another. The anterior and posterior margins of the costal are developed parallel to each other on the preserved part.

UU PL 704 (Fig. 3C): This specimen corresponds to a fragment of a costal, as is shown by the presence of the rib on the visceral part. Its morphology is similar with the previously described specimen (UU PL 703), but a larger part of the costal is preserved.

NHMC 21.7.3.1671 (Fig. 3D): This specimen most probably corresponds to a process of the plastron. It is long and flattened, with an elliptical cross-section. Further identification is not possible.

Remarks: The available specimens can be attributed to Pan-Trionychidae based on the presence of sculpturing that covers all metaplastic portions of the shell bones (Vitek and Joyce, 2015). The preserved pan-trionychid material from Plakias consists mainly of carapace fragments and a possible plastron element. Given that the sculpturing pattern is highly variable among soft-shelled turtles, even within individuals of the same species (Gardner and Russell, 1994; Vitek and Joyce, 2015), further identification is not possible. Compared to the only other known fossil pan-trionychid from Greece from the Pliocene of Gefira, northern Greece (Vlachos et al., 2015a), the Plakias specimens pertain to a smaller sized form, both in regards of carapacial disk length and thickness of the carapace. Another difference between these two Greek occurrences is noted in the sculpturing of the two forms: on the basis of the commonly preserved distal part of the costal, the pits on the Plakias pan-trionychid are mainly separated, whereas in the Gefira pan-trionychid, they are mostly coalesced, forming continuous grooves. As was mentioned above, however, the extreme variability of sculpturing that is observed within pan-trionychids does not allow us to determine taxonomic differences between the two Greek forms on the basis of this character.

GEOEMYDIDAE Theobald, 1868

196 *Mauremys* Gray, 1869

197 *Mauremys* sp.

198 **Referred specimens:** NHMC 21.7.3.1618, part of anterior lobe; NHMC 21.7.3.1619, right

199 xiphiplastron; NHMC 21.7.3.1620, left hypoplastron; NHMC 21.7.3.1621, group of 15

200 plastral fragments; NHMC 21.7.3.1622, neural; NHMC 21.7.3.1623, neural; NHMC

201 21.7.3.1624, neural; NHMC 21.7.3.1625, left hyoplastron fragment; NHMC 21.7.3.1626, left

202 hypoplastron fragment; NHMC 21.7.3.1627, right hypoplastron fragment; NHMC

203 21.7.3.1628, costal fragment; NHMC 21.7.3.1629, right hypoplastron fragment; NHMC

204 21.7.3.1630, costal fragment; NHMC 21.7.3.1631, costal fragment; NHMC 21.7.3.1632, left

205 epiplastron; NHMC 21.7.3.1633, left hyoplastron fragment; NHMC 21.7.3.1634, group of 23

206 shell fragments; NHMC 21.7.3.1635, right epiplastron; NHMC 21.7.3.1636, left epiplastron;

207 NHMC 21.7.3.1637, left epiplastron; NHMC 21.7.3.1638, costal fragment; NHMC

208 21.7.3.1639, costal fragment; NHMC 21.7.3.1640, two costals in association; NHMC

209 21.7.3.1641, right peripheral 1; NHMC 21.7.3.1642, right xiphiplastron fragment; NHMC

210 21.7.3.1643, right hypoplastron fragment; NHMC 21.7.3.1644, left hyoplastron fragment;

211 NHMC 21.7.3.1645, costal fragment; NHMC 21.7.3.1646, costal fragment; NHMC

212 21.7.3.1647, costal fragment; NHMC 21.7.3.1648, group of approximately 80 shell

213 fragments; NHMC 21.7.3.1649, neural fragment; NHMC 21.7.3.1650, group of 10 shell

214 fragments; NHMC 21.7.3.1651, neural I; NHMC 21.7.3.1652, costal fragment; NHMC

215 21.7.3.1653, costal fragment; NHMC 21.7.3.1654, costal fragment; NHMC 21.7.3.1655,

216 costal fragment; NHMC 21.7.3.1656, costal fragment; NHMC 21.7.3.1657, costal fragment;

217 NHMC 21.7.3.1658, costal fragment; NHMC 21.7.3.1659, costal fragment; NHMC

218 21.7.3.1660, costal fragment; NHMC 21.7.3.1661, costal fragment; NHMC 21.7.3.1662,

219 costal fragment; NHMC 21.7.3.1663, costal fragment; NHMC 21.7.3.1664, costal fragment;

220 NHMC 21.7.3.1665, costal fragment; NHMC 21.7.3.1666, peripheral fragment; NHMC

221 21.7.3.1667, peripheral fragment; NHMC 21.7.3.1668, peripheral fragment; NHMC
222 21.7.3.1669, group of 50 shell fragments; UU PL 705, possible hyoplastron fragment.

223 **Description:** The material consists of numerous shell fragments of relatively good
224 preservation. Most specimens preserve sufficient anatomical information, but several remain
225 indeterminate. The description of the most complete specimens is given below.

226

227 Carapace elements (Figs. 4A–J):

228 NHMC 21.7.3.1623 (Fig. 4A): This specimen corresponds to a complete neural. It is
229 hexagonal, with short anterior lateral sides. It is not crossed by any vertebral sulci, suggesting
230 that it is the second or fourth neural. NHMC 21.7.3.1645 (Fig. 4C) shows a similar
231 morphology.

232 NHMC 21.7.3.1622 (Fig. 4B): This is a complete neural that is hexagonal, with short
233 lateral sides. It is crossed by the vertebral sulci in the posterior part, suggesting that it is the
234 third or fifth neural.

235 NHMC 21.7.3.1624 (Fig. 4D): This complete neural is quadrangular to rounded, with
236 the anterior part being slightly wider. It is not crossed by any vertebral sulci. Its size is rather
237 small, in comparison to the other neurals described. A dorsal keel is noted longitudinally,
238 suggesting that it could belong to a young individual.

239 NHMC 21.7.3.1651 (Fig. 4E): This specimen corresponds to a complete first neural. It
240 is quadrangular with rounded edges, being longer than wide. The posterior part is crossed by
241 the vertebral sulcus that is not straight.

242 NHMC 21.7.3.1649 (Fig. 4F): This specimen corresponds to a fragment of a neural. It
243 is much wider than long, hexagonal in shape, with shorter anterior lateral sides. As such, it is
244 most probably one of the posterior neurals. Viscerally, the attachment for the vertebra is
245 visible. As it is not crossed by any sulci, it could be either the sixth or seventh neural.

NHMC 21.7.3.1640 (Fig. 4G): This specimen corresponds to the medial parts of two successive left costals in association. Both show the long and short medial sutures for the corresponding hexagonal neurals. In the preserved part, the sulci between the vertebrae and the pleurals can be observed.

NHMC 21.7.3.1638 (Fig. 4H): This specimen corresponds to an almost complete right costal, missing only the distal part. In medial side, two sutured surfaces are visible, one long and the other short, corresponding to an hexagonal neural. Dorsally, the sulci between the vertebrae are visible, suggesting that this could be the third or fifth costal.

NHMC 21.7.3.1639 (Fig. 4I): This specimen corresponds to the medial part of a left costal. In medial side, two sutured surfaces are visible, one long and one short, corresponding to an hexagonal neural. Dorsally, the sulci between the pleurals cross the medial part of this costal, suggesting that this could be the second, fourth or sixth costal.

NHMC 21.7.3.1641 (Fig. 4J): This specimen corresponds to an almost complete right peripheral I. Based on the preserved anterior border we can estimate the presence of a wide nuchal notch affecting also the first peripherals. The peripheral is long and narrow. Medially, the vertebral I and the pleural I show a long overlap on the peripheral, whereas the vertebral I contacts marginal II.

Plastron elements (Figs. 4K–S)

NHMC 21.7.3.1618 (Fig. 4K): This specimen corresponds to the right part of the anterior lobe of the plastron, consisting of the right epiplastron and most of the entoplastron. The epiplastron is long and narrow. Viscerally, a long but shallow lip is formed, being concave medially and convex laterally. Anteriorly, a shallow notch is noted. The entoplastron is hexagonal and rounded, being wider posteriorly. The gular scutes are wide and long, overlapping the anterior part of the entoplastron. The gular / humeral sulcus is slightly convex

laterally, and causes a slight constriction in the anterior part of the lobe. The humerals are medially short and laterally longer. The entoplastron is also overlapped by the anterior part of the pectorals. The humero-pectoral sulcus is slightly concave medially.

NHMC 21.7.3.1637 (Fig. 4L): This specimen corresponds to a left epiplastron. The epiplastron is long and rather wide. Viscerally, a long lip is formed, being concave medially and convex laterally. The gularscutes are wide and long, overlapping the anterior part of the entoplastron. The gularo / humeral sulcus is slightly convex laterally. The left epiplastron NHMC 21.7.3.1636 shows a similar morphology.

NHMC 21.7.3.1635 (Fig. 4M): This specimen corresponds to a right epiplastron. It is long and rather wide. Viscerally, a long and shallow lip is formed, being concave medially and convex laterally. The gularscutes are wide and long, overlapping the anterior part of the entoplastron. The gularo / humeral sulcus is slightly convex laterally and causes a slight constriction in the anterior part of the lobe.

NHMC 21.7.3.1632 (Fig. 4N): This is an almost complete left epiplastron, rather long and wide. Viscerally, a short, shallow and slightly concave lip is formed. It is covered medially by the gulars. An irregular growth of a small scute is noted in the anterior end of the gularo / humeral sulcus.

NHMC 21.7.3.1620 (Fig. 4O): This specimen corresponds to an almost complete left hypoplastron. The anterior suture, which connects it to the hyoplastron, is straight, whereas the posterior one, which connects it to the xiphiplastron, is slightly convex. The hypoplastron is rather flat. Viscerally, the abdominal forms a wide and slightly convex lip on the posterior lobe. Ventrally, an unusual morphology is noticed. Although the posterior part of the hypoplastron is covered by the abdominal scute, anteriorly there is another sulcus. Such sulcus is apparent also in another specimen from Plakias (NHMC 21.7.3.1643: fig. 4P), a morphology that has not been previously noted in geoemydids. However, the absence of

corresponding hyoplastra and the disarticulated nature of the material do not allow us to clarify this character.

NHMC 21.7.3.1643 (Fig. 4P): This specimen corresponds to an almost complete right hypoplastron. The anterior suture, which connects it to the hyoplastron, is straight, whereas the posterior one, which connects it to the xiphiplastron, is slightly convex. The hypoplastron is rather flat. Viscerally, the abdominal forms a wide and slightly convex lip on the posterior lobe. In the posterior part, the abdomino-femoral sulcus is clearly visible, whereas in the anterior part the sulcus is not clear as a result of erosion.

NHMC 21.7.3.1619 (Fig. 4Q): This specimen corresponds to an almost complete right xiphiplastron. The lateral sides of the posterior lobe are converging posteriorly. On the posterior part, a deep and wide anal notch is formed. The xiphiplastral extremities are rounded. The femorals cover the anterior part of the xiphiplastron, whereas the anals show a somewhat long covering on the posterior part of the lobe. The femoro-anal sulcus is slightly convex, being oriented antero-medially.

NHMC 21.7.3.1642 (Fig. 4R): This specimen corresponds to a fragment of the right xiphiplastron. On the basis of the preserved part, the presence of a wide angular anal notch can be estimated. In the anterior part, a short part of the femoro-anal sulcus is noted.

UU PL 705 (Fig. 4S): In the collections of UU, this small-sized specimen probably represents a fragment of the left hyoplastron. It preserves a curved sulcus that could be identified as the pectoro-abdominal one. Further identification is not possible, but it is overall similar with the respective material from NHMC and can be attributed to the same taxon as well.

Remarks: The specimens can be attributed to Geoemydidae on the basis of the following characters: first neural quadrangular, remaining neurals hexagonal with short anterolateral sides, pectorals medially shorter than the gulars, and deep anal notch. They can be further

321 identified as a member of *Mauremys*, on the basis of the contact between vertebral I with
322 marginal II and the entoplastron being crossed by the gulars and the humero-pectoral sulcus
323 (Claude et al., 2007). The *Mauremys* material from Plakias suggests attribution to a single
324 taxon, on the basis of the similar size, surface sculpturing and overall morphology. The
325 combined information of the preserved elements allows the documentation of the morphology
326 of most parts of the shell, indicating a confident assignment to *Mauremys*, but no species
327 determination can be made with certainty. Based on the available epiplastra, at least three
328 adult individuals are preserved, and a juvenile individual is also present. The unusual
329 morphology noted in the hypoplastra (NHMC 21.7.3.1620 and NHMC 21.7.3.1643) needs to
330 be further investigated, as it has not previously been observed in other geoemydids. However,
331 the absence of corresponding hyoplastra and the disarticulated nature of the material do not
332 allow us to interpret this character. The extended covering of the pleural I and vertebral I on
333 peripheral I distinguishes the Plakias *Mauremys* from the extant *Mauremys caspica* (Gmelin,
334 1774) and *Mauremys rivulata* (Valenciennes, 1833), as also from the extinct *Mauremys*
335 *gaudryi* (Depéret, 1885) (Pliocene, France; Hervet, 2003). As such, the Plakias *Mauremys* is
336 more similar to other Miocene terrapins from central and eastern Mediterranean (e.g.
337 *Mauremys campanii* Chesi et al., 2009, from Tuscany, Italy). The narrower neural I, the shape
338 of the entoplastron, having the posterior part shorter than the anterior one, and the shorter
339 pectoral covering on the entoplastron differentiate the Plakias geoemydid from *M. campanii*.
340 The angular anal notch also differentiates the Plakias geoemydid from the roughly similar
341 *Mauremys sarmatica* (Purschke, 1885) from the Miocene of Germany, which has a rounded
342 anal notch (Hervet, 2003). The Plakias terrapin represents the oldest described occurrence of
343 Geoemydidae from Greece, being older than the Allatini (Miocene / Pliocene boundary)
344 (Vlachos et al., 2015b) and the Maramena (latest Miocene) forms (Gad, 1990; Georgalis and
345 Kear, 2013; Vlachos et al., 2015b), both recovered from northern Greece. All other fossil

346 geoemydid occurrences from Greece are only known from Pleistocene and Holocene remains
347 (Chesi et al., 2007; Georgalis and Kear, 2013).

348

349 CROCODYLIA Gmelin, 1789

350 Crocodylia indet.

351 **Referred specimens:** UU PL 706 - UU PL 732, 27 isolated teeth; UU PL 735 - UU PL 736,
352 two phalanges.

353 **Description:** UU PL 706 - UU PL 732 (Fig. 5): Several isolated teeth preserve only a crown
354 that is characterized by being conical, variably pointed apically, and regularly concave
355 basally. Some of the teeth are only partially preserved; the largest crown is 8.3 mm long.
356 Mesiodistal carinae separate a lingual, slightly concave surface from a labial, slightly convex
357 surface. The carinae are not serrated. Both lingual and labial surfaces can be slightly wrinkled
358 and bear longitudinal ridges. Some of the teeth are slender, long, and pointed, whereas others
359 are more massive and apically blunt.

360 UU PL 735 - UU PL 736: These two phalanges are elongated elements showing a
361 single roundish articular surface proximally. The best preserved element (UU PL 735) is 18
362 mm long. Its distal portion is dorsoventrally flattened and laterally provided, on both sides,
363 with a sort of weak keel.

364 **Remarks:** The morphology of the teeth is fully congruent with that of generalized
365 crocodylians, but does not allow a more precise identification, since in most cases crocodylian
366 teeth are not diagnostic. These are, however, congruent with the morphology of the
367 *Crocodylus* teeth from the late Miocene of Italy (Delfino et al., 2007). Not much can be said
368 about the phalanges, except for the fact that they show standard crocodylian morphology.
369 They are referred to the same taxon as the teeth. The Plakias specimens constitute the first

370 fossils of crocodylians described from Greece, as also one of the last occurrences of this
371 group in the European continent (Table 1).

372

373 SQUAMATA Oppel, 1811

374 AMPHISBAENIA Gray, 1844

375 *Amphisbaenia* indet.

376 **Referred specimen:** UU PL 733, a single presacral vertebra.

377 **Description:** The specimen is a rather small, procoelous vertebra, with a centrum length of
378 only 1.2 mm (Fig. 6). A slight degree of deformation is recognizable in anterior view. The
379 centrum is dorsoventrally compressed and has a flattened ventral surface and subparallel
380 lateral margins. A massive and rounded synapophysis is visible on the right side of the
381 vertebra. In dorsal view, the neural arch is constricted in the middle. Its dorsal surface is
382 flattened and the neural spine is lacking, as well as the zygosphenes. Only the right
383 prezygapophysis is preserved: it is roughly sub-elliptical and tilted dorsally about 30°. The
384 prezygapophyseal process is rather short. Neither the posterior end of the neural arch nor the
385 postzygapophyses are preserved.

386 **Remarks:** The specimen can be attributed to *Amphisbaenia* on the basis of the combination
387 of the following characters: small size, dorsoventrally compressed centrum with a flattened
388 ventral surface and roughly parallel lateral margins, short and robust prezygapophyses,
389 massive and rounded synapophyses, absence of zygosphenes, and a dorsally flattened neural
390 arch lacking a neural spine (Estes, 1983; Delfino, 2003). *Amphisbaenians* are present in the
391 extant herpetofauna of Greece, with *Blanus* occurring in the Dodecanese Islands (see below in
392 Biogeography), but they were totally absent in the fossil record of the country. As such, the
393 specimen described herein represents the first known fossil *amphisbaenian* from Greece. Its
394 affinities with *Blanidae*, which are the only extant *amphisbaenians* inhabiting Europe, cannot

be tested on the basis of the vertebral morphology: isolated vertebrae of *Amphisbaenia* do not show significant diagnostic features that would allow for a more precise identification (Estes, 1983). The sole other known fossil amphisbaenian from the Aegean region has been recovered from the Pliocene of Çalta, Turkey (Rage and Sen, 1976). The Çalta amphisbaenian is represented by three vertebrae, none of which was figured or described in detail. As a matter of fact, no further comparison between these specimens can be made.

Interestingly also, the vertebra UU PL 733 represents the sole non-snake squamate from Plakias, as other lizards are conspicuously absent. We consider that this absence is due to taphonomic or collection biases and does not reflect the actual lizard palaeodiversity of Plakias.

SERPENTES Linnaeus, 1758

COLUBROIDEA Oppel, 1811

?NATRICINAE Bonaparte, 1838 (sensu Szyndlar, 1991b)

?Natricinae indet.

Referred specimen: UU PL 734, a single precloacal vertebra.

Description: The vertebra is fragmentary, with the prezygapophyses, part of the cotyle and part of the neural spine being eroded (Fig. 7). The vertebra is procoelous, relatively large and proportionally elongate, with a centrum length of 5.5 mm. The prezygapophyseal processes are not clearly visible, due to the fact that the anterior portion of the vertebra is not well preserved. The cotyle is rather incomplete but appears to be relatively rounded. Only part of the zygosphenes is preserved. The condyle is spherical and rather robust, and protrudes significantly from the posterior part of the vertebra. The roof of the zygantrum is visible and appears to be relatively thick. The neural canal is rather large, almost equal in size to the condyle. Synapophyses are divided in diapophyses and parapophyses. The neural spine is

broken. The hypapophysis was probably present, however, this cannot be evaluated with certainty. In ventral view, a subcentral foramen is also visible at mid centrum length of the vertebra.

Remarks: The specimen can be attributed to *Serpentes* on the basis of the general vertebral shape and especially the presence of zygantrum and zygosphenes (Rage, 1984). The single snake vertebra from Plakias is fragmentary, thus precluding the evaluation of exact taxonomic affinities. However, the probable presence of hypapophysis could indicate that this specimen could belong to either natricines, viperids or elapids. All these three groups have been recorded in the Miocene of Greece (Szyndlar, 1991a, b, 1995; Szyndlar and Rage, 2002; Georgalis et al., 2016a). The combination, however, of a rather elongate centrum, strong subcentral ridge and, probably, a posteriorly vaulted neural spine, prompts us to consider natricine affinities as the most plausible for the Plakias snake. This identification is further supported by direct comparison with numerous skeletons of extant natricine snakes.

5. Discussion

5.1 Palaeoecology of *Plakias*

The locality of Plakias is mostly known for its micromammal assemblage (de Bruijn and Meulenkamp, 1972; de Bruijn et al., 2012). This is comprised of erinaceomorphs and soricomorphs eulipotyphlans, and eomyid, sciurid, glirid and murid rodents, whereas larger mammals are known only by an indeterminate suid similar to *Propotamochoerus* (van der Made, 1996; Koufos, 2006; de Bruijn et al., 2012). The presence of at least three distinct taxa of sciurids (Koufos, 2006) indicates the likely presence of a forested environment (de Bruijn et al., 1980), although we acknowledge that several fossil and extant sciurids are ground dwellers (e.g. Viriot et al., 2011). Plant remains and invertebrates are also known from Plakias (de Bruijn et al., 2012). Fossil invertebrates include fresh-water gastropods, such as

Planorbis and *Brotia* (de Bruijn et al., 2012), which further suggest the presence of a river or lake system.

The new amphibian and reptile finds from Plakias allow us to draw further conclusions regarding the palaeoecology of this locality. The presence of an alytid frog implies wet habitats, which is the common environment for the extant members of the group (Vitt and Caldwell, 2014). All extant Crocodylia are known to be water dwellers, inhabiting river systems, lakes, and even venture into the open sea (Steel, 1973; Britton et al., 2012). The fragmentary nature of the Cretan crocodylian does not allow us to make a proper specific or generic identification, but as the teeth from Plakias bear strong resemblance with certain Neogene finds from the Mediterranean that are assigned to Crocodyloidea or Alligatoroidea, we can infer a similar lifestyle. All extant pan-trionychids are known to be strictly aquatic, and a similar or identical life strategy has been proposed for all fossil taxa of this group as well (Vitek and Joyce, 2015 and references therein). The presence of a geoemydid further indicates the presence of lake and river systems (Busack and Ernst, 1980). The amphisbaenian vertebra is fragmentary and cannot be assigned to the specific level. However, the vast majority of amphisbaenians are fossorial (Kearney, 2003) and as such, a similar, burrowing lifestyle is also proposed for the Cretan representative of this clade. Additionally, the presence of a natricine snake adds a further aquatic or semi-aquatic taxon to the locality (Vitt and Caldwell, 2014).

5.2 Biogeography

At least until the late Serravallian (middle Miocene), Crete was located at the southern part of Aegæis, the continental area that united modern Greece with Anatolia, but it was most probably already isolated as an island by the Tortonian (late Miocene), after the opening of the Proto-Aegean Sea (Dermitzakis and Papanikolaou, 1981; Poulakakis et al., 2005). The

suggested age of the Plakias fauna as early Tortonian (de Bruijn et al., 2012) coincides with the breakup of the southern Aegean landmass and the early formation of Crete as an island (Poulakakis et al., 2005: fig. 4). As such, we cannot determine with certainty whether Plakias hosted a truly continental fauna or an insular fauna at least partly deriving from the former continental assemblages. However, even if the fauna was an insular one, it should have strong biogeographic affinities with coeval mainland Aegäis faunas.

The micromammal fauna of Plakias bears strong affinities with coeval ones from Central Europe (de Bruijn et al., 2012). The amphibian and reptile fossils described herein provide additional information about the palaeobiogeography of this part of southeastern Europe, as Miocene herpetofaunas are not well documented in that region (Georgalis et al., 2013; Georgalis and Kear, 2013).

The presence of an alytid frog in the late Miocene of Crete is not of biogeographic importance, as this group was widespread throughout Europe during that time interval and has been also found from several other Neogene localities from Greece (Rage and Roček, 2003). All other fossil amphibians from Crete are of Quaternary age (Caloi et al., 1986; Sanchiz, 1998).

The two distinct turtle taxa recovered from Plakias represent the oldest turtles from Crete, as also the oldest occurrences of Geoemydidae and Pan-Trionychidae from Greece up to date. Turtles, in general, were considered up to now to be absent from Neogene localities of Crete, with their only remains known from Pleistocene and Holocene sediments (Bachmayer et al., 1975; Kotsakis, 1977; Brinkerink, 1996; Chesi et al., 2007; Georgalis and Kear, 2013). Although Brinkerink (1996: p. 208) noted that no turtles are known from pre-Pleistocene sites from Crete, he pointed out the potential of the site of Plakias. The presence of pan-trionychids in the early late Miocene of Greece represents the oldest record and the second only occurrence of this group from the country, which was otherwise only known from the

Pliocene of northern Greece (Vlachos et al., 2015a). Pan-trionychids are widespread throughout Europe during the Neogene (Hummel, 1932; Młynarski, 1976; Danilov, 2005; Karl, 1999), however, their alpha taxonomy has not been settled with certainty, mostly because of the homoplastic characters that are observed in their morphology (Vitek and Joyce, 2015). Curiously also, Miocene pan-trionychids were totally absent until now from Greece, despite the wealth of fossil localities of that age in the country and extensive sampling (Georgalis and Kear, 2013). The new specimens from Plakias could probably indicate that this absence was not genuine and that pan-trionychids will be subsequently found in other Greek Miocene localities. This is further supported by the presence of pan-trionychids in the Miocene of Bulgaria (Pamouktchiev et al., 1998; Georgalis and Kear, 2013) and Anatolia (Staesche et al., 2007). Affinities of the new Cretan pan-trionychid with the Bulgarian and Anatolian forms cannot be established due to the fragmentary nature of the new specimens. Other, relatively geographically close Miocene pan-trionychids are also known from Cyprus (Hadjisterkotis et al., 2000), Egypt (Lapparent de Broin, 2000), Italy and Malta (Kotsakis, 1985). The Plakias geoemydid adds to the Miocene diversity of this group in southeastern Europe. Additionally, this pre-Messinian occurrence of *Mauremys* further corroborates the suggestion of Chesi et al. (2009) for warm and wet conditions during that period in southern Mediterranean Europe.

Crocodylians are conspicuous elements in several Miocene faunas across European localities (among others, Ginsburg and Bulot, 1997; Kotsakis et al., 2004; Delfino et al., 2007; Delfino and Rook, 2008; Martin, 2010; Martin and Gross, 2011; Delfino and Rossi, 2013). However, they are relatively scarce by the late Miocene and were apparently absent from southeastern Europe and Anatolia (Böhme, 2003; Sen et al., 2011). The youngest up to date published records of crocodylians from Europe are known from the latest Miocene of Italy (Table 1), as supposed occurrences from the late Miocene and Pliocene of Spain and Portugal

have not been accompanied by descriptions or figures, and are here considered anecdotal (Delfino et al., 2007). The taxon *Diplocynodon levantinum* from Bulgaria, was initially thought to be of Pliocene age (Huene and Nikoloff, 1963), whereas it was later treated questionably as late Miocene (Delfino and Rossi, 2013), but its type locality is now believed to pertain most probably to the middle Miocene (Sen et al., 2011). With the exception of the slender snouted forms attributed to *Tomistoma* (Capellini, 1890; Vianna and Moraes, 1945), several late Miocene occurrences are considered to be members of *Crocodylus* or at least cf. *Crocodylus* (Delfino et al., 2007; Delfino and Rook, 2008; Delfino and Rossi, 2013) that possibly dispersed from Africa, well before the so-called Messinian Salinity Crisis (Delfino et al., 2007). Interestingly also, Gargano and Scontrone, that yielded *Crocodylus* remains, were palaeoislands during the latest Miocene; we can thus speculate that the Cretan crocodylian could also belong to the same lineage and had originated through a similar dispersal route from Africa. Whatever the case, the isolated teeth from Plakias represent the first crocodylians from Greece to be formally described and one of only few late Miocene occurrences of this group in Europe.

Our knowledge of squamate biogeography during the Miocene of Europe is hindered by the lack of consensus surrounding the taxonomy and phylogenetic relationships of its representatives. This is especially true for the southeastern European localities, from where only few specimens have been recovered, most of which originating from classic, well known mammal-bearing sites (Gaudry, 1862-67; Weithofer, 1888; Richter, 1995; Georgalis et al., 2016b). The presence of a natricine in the late Miocene of Plakias adds to the already known diversity of this widespread snake group (Szyndlar, 1991b), and in fact, represents one of the southernmost fossil occurrences of colubroids in Europe. Whether the Plakias snake bears close affinities with other Neogene natricines from southeastern Europe (e.g. the natricine from Maramena described by Szyndlar [1995]) cannot be tested due to the fragmentary nature

545 of the single known specimen. Furthermore, the *Plakias* natricine represents the sole Neogene
546 record of snakes from Crete, as all other fossil occurrences from the island are confined to
547 Pleistocene and Holocene sediments (Holman, 1998). Amphisbaenians are well known from
548 several Miocene European localities, but almost only from the western and central parts of the
549 continent (Roček, 1984; Bolet et al., 2014; Čerňanský et al., 2015), whereas they have never
550 been described from the Balkan Peninsula and eastern Europe in general (Delfino, 2003), with
551 the single exception of a blandid from the middle Miocene of Tauț, Romania (Venczel and
552 Știucă, 2008). After the end of the Miocene, the distribution of amphisbaenians in Europe
553 became gradually restricted to its Mediterranean margins, becoming extinct from most
554 regions after the Pleistocene and surviving today only in the Iberian Peninsula and few Greek
555 Islands (Delfino, 1997; Delfino and Bailon, 2000). Amphisbaenians are represented in the
556 Greek extant herpetofauna solely by the species *Blanus strauchi* (Bedriaga, 1884), which is
557 distributed in the islands of Samos, Fournoi, Leros, Kos, Symi, Rhodes and Kastellorizon
558 (Valakos et al., 2008). *Blanus* occurs also in the adjacent Anatolia, represented by *B. strauchi*
559 and two additional, recently described or revalidated species: *Blanus alexandri* Sindaco,
560 Kornilios, Sacchi and Lymberakis, 2014, and *Blanus aporus* Werner, 1898. The Anatolian
561 fossil record of amphisbaenians is also poor, consisting of only a single record from the
562 Pliocene of Çalta (Rage and Sen, 1976; Delfino, 1997). Whether or not the *Plakias* specimen
563 represents a species of *Blanus*, cannot be evaluated on the basis of vertebral characters. It is,
564 however, probable that the *Plakias* amphisbaenian belongs indeed to *Blanus*, as members of
565 this genus have a wide Miocene distribution in western and Central Europe (Bolet et al.,
566 2014; Čerňanský et al., 2015) and the disjunct extant restricted distribution in the western and
567 eastern edges of the continent has been in fact interpreted as relics of an once continuous
568 range in southern Europe (Alexander, 1966; Delfino, 2003), a situation that has also been
569 observed for other squamate groups, such as erycid and elapid snakes (Szyndlar, 1991a, b;

Rage, 2013). In fact, Greece and in general the southern Balkans have been suggested as acting as a refugium for multiple squamate groups, such as scolecophidians, erycids, the large anguid *Pseudopus* and the so called “Oriental Vipers” (Delfino, 2003), which all have their sole extant European populations only in that region (Sindaco and Jeremčenko, 2008). However, an alternative hypothesis of the Plakias amphisbaenian sharing affinities with African ones should also be taken into consideration, due to the geographical proximity of Crete to Africa and the unexpected but now well known marine dispersal capabilities of these squamates (Kearney, 2003; Longrich et al., 2015), but such hypothesis is severely hindered by the poor fossil record of African squamates (Rage, 2003; Delfino et al., 2004). Pending the discovery of more complete material from Plakias that could ideally comprise cranial elements that bear diagnostic features, no further biogeographic correlations of the Cretan amphisbaenian with European, Anatolian or African taxa can be made with certainty.

6. Conclusions

The herpetofauna of Plakias described in this paper includes the oldest amphibians and reptiles from the Island of Crete and represents one of the southernmost fossil herpetofaunas of Europe. The alytid frog presented herein adds to the known record of this group from the Miocene of southern Europe. The crocodylians are the first such described faunal elements from the country and represent one of the youngest occurrences of this group in Europe. Turtles include two distinct taxa: a geoemydid and a pan-trionychid, both representing the oldest occurrences of these groups from Greece. The single colubroid snake specimen adds further to the published record of Miocene snakes from Greece, whereas the amphisbaenian vertebra from Plakias represents the first described fossil of this group from the country, suggesting that amphisbaenians had a continuous range in the northern Mediterranean area. The herpetofauna of Plakias is shown to be diverse. Overall, it further adds to our knowledge

of the Miocene herpetofaunas of southeastern Europe, a region in which amphibian and reptile fossils are still not adequately known.

Acknowledgements

Wilma Wessels and Hans de Bruijn (University of Utrecht) are thanked for the long term loan of the material utilized in this study. Moysis Mylonas and Charalampos Fassoulas (Natural History Museum of Crete and University of Crete) provided access to the turtle material to EV. We also thank Walter Joyce (University of Fribourg), Borja Sanchiz (Museum of Natural Sciences, Madrid), George Iliopoulos (University of Patras), Dimitris Kostopoulos (Aristotle University of Thessaloniki) and George Koufos (Aristotle University of Thessaloniki), as also our reviewers Stefan Vasile (University of Bucharest), Johannes Müller (Museum für Naturkunde, Berlin) and Jean-Claude Rage (Muséum National d'Histoire Naturelle, Paris), and editors Gilles Escarguel (University of Lyon), and Gildas Merceron (University of Poitiers), for valuable comments that improved the quality of the manuscript. Study of snake skeletons that were used for comparisons in this study was funded by SYNTHESYS ES-TAF-5910 (Museo Nacional de Ciencias Naturales, Madrid) and SYNTHESYS AT-TAF-5911 (Naturhistorisches Museum, Vienna) grants to GLG, and the curators of these institutions, respectively Marta Calvo-Revuelta and Heinz Grillitsch, are also acknowledged here. GLG acknowledges travel support from the University of Fribourg. Work is supported by Fondi di Ateneo dell'Università di Torino (2013-2014) and Generalitat de Catalunya (2014 SGR 416 GRC).

References

Abbazzi, L., Delfino, M., Gallai, G., Trebini, L., Rook, L., 2008. New data on the vertebrate assemblage of Fiume Santo (North-western Sardinia, Italy), and overview on the Late Miocene Tusco-Sardinia paleobioprovince. *Palaeontology* 51, 425–451.

- 621 Alexander, A.A., 1966. Taxonomy and variation of *Blanus strauchi* (Amphisbaenia, Reptilia).
 622 Copeia 1966, 205–224.
- 623 Antunes, M.T., 1994. On Western Europe Miocene gavials (Crocodylia): their
 624 paleogeography, migrations and climatic significance. Comunicações do Instituto
 625 Geológico e Mineiro 80, 57–69.
- 626 Augé, M., 2005. Evolution des lézards du Paléogène en Europe. Publications Scientifiques du
 627 Muséum, Mémoires du Muséum national d'histoire naturelle, Paris 192, 1–369.
- 628 Bachmayer, F., 1967. Eine Riesenschildkröte aus den altplozänen Schichten von Pikermi
 629 (Griechenland). Annales Géologiques des Pays Helléniques 19, 512–526.
- 630 Bachmayer, F., Brinkerink, J., Symeonidis, N., 1975. Pleistozäne Schildkröten aus Höhlen der
 631 Insel Kreta. Annales Géologiques des Pays Helléniques 27, 110–121.
- 632 Bailon, S., 1999. Différenciation ostéologique des anoures (Amphibia, Anura) de France.
 633 Fiches d'osteologie animale pour l'archéologie (C: Varia) 1, 1–41.
- 634 Batsch, A.J.G.C., 1788. Versuch einer Anleitung, zur Kenntniß und Geschichte der Thiere
 635 und Mineralien, für akademische Vorlesungen entworfen, und mit den nöthigsten
 636 Abbildungen versehen. Erster Theil. Allgemeine Geschichte der Natur; besondre der
 637 Säugthiere, Vögel, Amphibien und Fische. Akademische Buchhandlung, Jena.
- 638 Bedriaga J. 1884 Nachträgliche Bemerkung über *Amphisbaena strauchi* v. Bedr. Zoologischer
 639 Anzeiger 7, 346.
- 640 ~~Boettger, O., 1883. Herpetologische Mittheilungen. Berichte des Offenbacher Vereins für~~
 641 ~~Naturkunde (1880–1882) 22–23, 147–156.~~
- 642 Böhme, M., 2003. The Miocene Climatic Optimum: evidence from ectothermic vertebrates of
 643 Central Europe. Palaeogeography, Palaeoclimatology, Palaeoecology 195, 389–401.

644 Bolet, A., Delfino, M., Fortuny, J., Almécija, S., Robles, J.M., Alba, D.M., 2014. An
 645 amphisbaenian skull from the European Miocene and the evolution of Mediterranean
 646 worm lizards. PLoS One 9, e98082.

647 Bonaparte, C.L. 1832–1841. Iconographia della fauna Italica per le quattro classi degli
 648 animali vertebrati. 2. Anfibi. Tipografia Salviucci, Roma.

649 Brinkerink, H.P., 1996. Tortoises and turtles from the Pleistocene of Crete, in: Reese, D.S.
 650 (Ed.), Pleistocene and Holocene fauna of Crete and its first settlers. Monographs in
 651 World Archaeology 28, Prehistory Press, 207–210.

652 Britton, A.R.C., Whitaker, R., Whitaker, N., 2012. Here be a Dragon: exceptional size in a
 653 saltwater crocodile (*Crocodylus porosus*) from the Philippines. Herpetological Review
 654 43, 541–546.

655 Brochu, C.A., 2000. Phylogenetic relationships and divergence timing of *Crocodylus* based
 656 on morphology and the fossil record. Copeia 2000, 657–673.

657 Busack, S.D., Ernst, C.H., 1980. Variation in Mediterranean populations of *Mauremys* Gray
 658 1869 (Reptilia, Testudines, Emydidae). Annals of the Carnegie Museum 49, 251–264.

659 Caloi, L., Kotsakis, T., Palombo, M.R., 1986. La fauna a vertebrati terrestri del Pleistocene
 660 delle isole del Mediterraneo. Geologia Romana 25, 235–256.

661 Capellini, G., 1890. Sul coccodrilliano garialoide (*Tomistoma calaritanus*) scoperto nella
 662 collina di Cagliari nel 1868. Rendiconti Reale Accademia Lincei (ser. 4) 6 (I sem.),
 663 149–151.

664 Čerňanský, A., Rage, J.-C., Klembara, J., 2015. The Early Miocene squamates of Amöneburg
 665 (Germany): the first stages of modern squamates in Europe. Journal of Systematic
 666 Palaeontology 13, 97–128.

667 Chesi, F., Delfino, M., Abbazzi, L., Carboni, S., Lecca, L., Rook, L., 2007. New fossil
 668 vertebrate remains from San Giovanni di Sinis (Late Pleistocene, Sardinia): the last

669 *Mauremys* (Reptilia, Testudines) in the central Mediterranean. Rivista Italiana di
 670 Paleontologia e Stratigrafia 113, 287–297.

671 Chesi, F., Delfino, M., Rook, L., 2009. Late Miocene *Mauremys* (Testudines, Geoemydidae)
 672 from Tuscany (Italy): evidence of terrapin persistence after a mammal turnover. Journal
 673 of Paleontology 83, 379–388.

674 Claude, J., Tong, H., Suteethorn, V., 2007. Turtles from the late Eocene–early Oligocene of
 675 the Krabi Basin (Thailand). Bulletin de la société géologique de France, Série 1, 178,
 676 305–316.

677 Danilov, I.G., 2005. Die fossilen Schildkröten Europas. Part 9, in: Fritz, U. (Ed.), Handbuch
 678 der Reptilien und Amphibien Europas. Part 3/IIIB: Schildkröten (Testudines) II. AULA-
 679 Verlag GmbH, Wiebeisheim, pp. 329–448.

680 Daza, J.D., Bauer, A.M., Snively, E., 2014. On the gekkotan fossil record. Anatomical Record
 681 297, 433–462.

682 de Bruijn H., Doukas, C.S., van den Hoek Ostende, L.W., Zachariasse, W.J., 2012. New finds
 683 of rodents and insectivores from the Upper Miocene at Plakias (Crete, Greece). Swiss
 684 Journal of Palaeontology 131, 61–75.

685 de Bruijn, H., Meulenkamp, J.E., 1972. Late Miocene rodents from the Pandanassa formation
 686 (Prov. Rethymnon), Crete, Greece. Proceedings of the Koninklijke Nederlandse
 687 Akademie van Wetenschappen, Series B 75, 54–60.

688 de Bruijn, H., van der Meulen, A.J., Katsikatsos, G., 1980. The mammals from the Lower
 689 Miocene of Aliveri (Island of Evia, Greece). Part 1: The Sciuridae. Proceedings of the
 690 Koninklijke Nederlandse Akademie van Wetenschappen, Series B 83, 241–261.

691 Delfino, M., 1997. *Blanus* from the early Pleistocene of southern Italy: another small tessera
 692 from a big mosaic, in: Böhme, I.W., Bischoff, W., Ziegler, T., (Eds.), Herpetologica
 693 Bonnensis. SEH, Bonn, pp. 89–97.

694 Delfino, M., 2003. A Pleistocene amphisbaenian from Sicily. *Amphibia-Reptilia* 24, 407–414.
 695 Delfino, M., Bailon, S., 2000. Early Pleistocene herpetofauna from Cava Dell' Erba and Cava
 696 Pirro (Apulia, Southern Italy). *Herpetological Journal* 10, 95–110.
 697 Delfino, M., Böhme, M., Rook, L., 2007. First European evidence for transcontinental
 698 dispersal of *Crocodylus* (late Neogene of southern Italy). *Zoological Journal of the*
 699 *Linnean Society, London* 149, 293–307.
 700 Delfino, M., Rook, L., 2008. African crocodylians in the late Neogene of Europe: a revision
 701 of *Crocodylus bambolii* Ristori, 1890. *Journal of Paleontology* 82, 336–343.
 702 Delfino, M., Rossi, M.A., 2013. Fossil crocodylid remains from Scontrone (Tortonian,
 703 Southern Italy) and the late Neogene Mediterranean biogeography of crocodylians.
 704 *Geobios* 46, 25–31.
 705 Delfino, M., Segid, A., Yosief, D., Shoshani, J., Rook, L., Libsekal, Y., 2004. Fossil reptiles
 706 from the Pleistocene *Homo*-bearing locality of Buia (Eritrea, Northern Danakil
 707 Depression), in: Abbate, E., Woldehaimanot, B., Libsekal, Y., Tecle, T.M., Rook, L.,
 708 (Eds.), A step towards human origins. The Buia *Homo* one-million-years ago in the
 709 Eritrean Danakil Depression (East Africa). *Rivista Italiana di Paleontologia e*
 710 *Stratigrafia* 110(supplement), 51–60.
 711 Depéret, C., 1885. Description géologique du bassin tertiaire du Roussillon. *Annales des*
 712 *Sciences Géologiques* 17, 1–272.
 713 Dermitzakis, D.M., Papanikolaou, D.J., 1981. Paleogeography and geodynamics of the
 714 Aegean region during the Neogene. *Annales Géologiques des Pays Helléniques* 30, 245–
 715 289.
 716 Estes, R., 1983. Sauria Terrestria, Amphisbaenia, in: Wellnhofer, P., (Ed.), *Encyclopedia of*
 717 *Paleoherpetology*, part 10a. G. Fischer Verlag, Stuttgart.

- 718 Fischer von Waldheim, G., 1813. Zoognosia. Tabulis Synopticis Illustrata, in: Usum
719 Praelectionum Academiae Imperialis Medico-Chirurgicae Mosquensis Edita. Ed. 3.
720 Volume 1. Nicolai Sergeidis Vsevolozsky, Moscow.
- 721 Fitzinger, L.J.F.J., 1843. Systema Reptilium. Fasciculus Primus. Braumüller et Seidel,
722 Vienna.
- 723 Frost, D.R., Grant, T., Faivovich, J., Bain, R.H., Haas, A., Haddad, C.F.B., de Sá, R.O.,
724 Channing, A., Wilkinson, M., Donnellan, S.C., Raxworthy, C.J., Campbell, J.A., Blotto,
725 B.L., Moler, P.E., Drewes, R.C., Nussbaum, R.A., Lynch, J.D., Green, D.M., Wheeler,
726 W.C., 2006. The amphibian tree of life. Bulletin of the American Museum of Natural
727 History 297, 1–370.
- 728 Gad, J., 1990. Nachweis einer Schildkröte des *Mauremys–Ocadia*-Komplexes aus dem
729 Obermiozän von Nordgriechenland. Salamandra 26, 311–313.
- 730 Garcia, G., Kostopoulos, D.S., Valentin, X., Koufos, G.D., 2016. Testudines, in: Koufos,
731 G.D., Kostopoulos, D.S., (Eds.), Palaeontology of the upper Miocene vertebrate
732 localities of Nikiti (Chalkidiki Peninsula, Macedonia, Greece). Geobios 49, 23–27.
- 733 Gardner, J.D., Russell, A.P., 1994. Carapacial variation among soft-shelled turtles
734 (Testudines: Trionychidae), and its relevance to taxonomic and systematic studies of
735 fossil taxa. Neues Jahrbuch für Geologie und Paläontologie–Abhandlungen 193, 209–
736 244.
- 737 Gaudry, A., 1862–1867. Animaux fossiles et géologie de l'Attique. Savy, Paris.
- 738 Gauthier, J.A., Kearney, M., Maisano, J.A., Rieppel, O., Behike, A.D.B., 2012. Assembling
739 the squamate tree of life: perspectives from the phenotype and the fossil record. Bulletin
740 of the Peabody Museum of Natural History 53, 3–308.
- 741 Georgalis, G.L., Kear, B.P., 2013. The fossil turtles of Greece: an overview of taxonomy and
742 distribution. Geobios 46, 299–311.

743 Georgalis, G.L., Szyndlar, Z., Kear, B.P., Delfino, M., 2016a. New material of *Laophis*
 744 *crotaloides*, an enigmatic giant snake from Greece, with an overview of the largest
 745 fossil European vipers. *Swiss Journal of Geosciences* 109, 103–116.

746 Georgalis, G.L., Velitzelos, E., Velitzelos, D.E., Kear, B.P., 2013. *Nostimochelone lampra*
 747 gen. et sp. nov., an enigmatic new podocnemidoidean turtle from the lower Miocene of
 748 northern Greece, in: Brinkman, D.B., Holroyd, P.A., Gardner, J.D., (Eds.), *Morphology*
 749 *and Evolution of Turtles*. Springer, Dordrecht, pp. 277–287.

750 Georgalis, G.L., Villa, A., Delfino, M., 2016b. First description of a fossil chamaeleonid from
 751 Greece and its relevance for the European biogeographic history of the group. *The*
 752 *Science of Nature* 103, 12.

753 Ginsburg, L., Bulot, C., 1997. Les *Diplocynodon* (Reptilia, Crocodylia) de l'Orléanien
 754 (Miocène inférieur à moyen) de France. *Geodiversitas* 19, 107–28.

755 Gmelin, J.F., 1789. Tom. I. Pars III, in: *Caroli a Linné, Systema Naturae*. Beer, G.E. Leipzig.
 756 pp. 1033–1516.

757 Gmelin, S.G., 1774. *Reise durch Rußland zur Untersuchung der drey Natur-Reiche*.
 758 Kaiserliche Akademie der Wissenschaften, St. Petersburg.

759 Gray, J.E., 1844. Catalogue of the tortoises, crocodiles, and amphisbaenians, in the collection
 760 of the British Museum. Trustees of the British Museum, London.

761 Gray, J.E., 1869. Description of *Mauremys laniaria*, a new freshwater tortoise. *Proceedings of*
 762 *the Zoological Society of London* 1869, 499–500.

763 Hadjisterkotis, E., Kassapis, I., Klerides, N., 2000. Paleontological evidence for the presence
 764 of *Trionyx* sp. on Cyprus, with new information on living and alien species of terrapins.
 765 Newsletter, Hellenic Zoological Society 33, 8–10.

766 Hervet, S., 2003. Le groupe "*Palaeochelys* sensu lato–*Mauremys*" dans le contexte
 767 systématique des Testudinoidea aquatiques du Tertiaire d'Europe occidentale: Apports à

768 la biostratigraphie et à la paléobiogéographie. Ph.D. thesis, Muséum National d' Histoire
 769 Naturelle, Paris.

770 Holman, A.J., 1998. Pleistocene amphibians and reptiles in Britain and Europe. Oxford
 771 Monographs on Geology and Geophysics 38, Oxford University Press, New York,
 772 Oxford.

773 Huene, F., von., Nikoloff, I., 1963. Ein pliozänes krokodil in Bulgarien. Neues Jahrbuch
 774 für Geologie und Paläontologie Abhandlungen 118, 266–271.

775 Hummel, K., 1932. Trionychia fossilia, in: Questedt, W., (Ed.), Fossilium Catalogus I:
 776 Animalia, Junk, W., Berlin.

777 Joyce, W.G., Parham, J.F., Gauthier, J.A., 2004. Developing a protocol for the conversion of
 778 rank-based taxon names to phylogenetically defined clade names, as exemplified by
 779 turtles. Journal of Paleontology 78, 989–1013.

780 Karl, H.V., 1999. Die Zoogeographie der känozoischen Weichschildkröte *Trionyxtriunguis*
 781 Forskål, 1775 (Testudines: Trionychidae). Joannea Geologie und Paläontologie, Graz 1,
 782 27–60.

783 Kearney, M., 2003. Systematics of the Amphisbaenia (Lepidosauria: Squamata) based on
 784 morphological evidence from recent and fossil forms. Herpetological Monographs 17,
 785 1–74.

786 Kotsakis, T., 1977. I resti di anfibi e rettili pleistocenici della grotta 'Bate' (Rethymnon,
 787 Creta). Atti della Accademia Nazionale dei Lincei 63, 571–582.

788 Kotsakis, T., 1985. Les Trionychidae (Testudinata, Reptilia) fossiles de l'Italie. Bollettino
 789 della Società Paleontologica Italiana 24, 161–168.

790 Kotsakis, T., Delfino, M., Piras, P., 2004. Italian Cenozoic crocodilians: taxa, timing and
 791 biogeographic implications. Palaeogeography, Palaeoclimatology, Palaeoecology 210,
 792 67–87.

793 Koufos, G.D., 2006. The Neogene mammal localities of Greece: faunas, chronology, and
794 biostratigraphy. *Annales Géologiques des Pays Helléniques* 4, 183–214.

795 Koufos, G.D., Kostopoulos, D.S., 2013. First report of *Brachypotherium* Roger, 1904
796 (Rhinocerotidae, Mammalia) in the Middle Miocene of Greece. *Geodiversitas* 35, 629–
797 641.

798 Koufos, G.D., Kostopoulos, D., Vlachou, T., 2005. Neogene/Quaternary mammalian
799 migrations in Eastern Mediterranean. *Belgian Journal of Zoology* 135, 181–190.

800 Lapparent de Broin, F., de, 2000. African chelonians from the Jurassic to present: phases of
801 development and preliminary catalogue of the fossil record. *Palaeontologia Africana* 36,
802 43–82.

803 Laurenti, J.N., 1768. *Specimen medicum, exhibens synopsin reptilium emendatam cum*
804 *experimentis circa venena et antidota reptilium austracorum, quod auctoritate et*
805 *consensu. Joan. Thomae, Vienna.*

806 Linnaeus, C., 1758. *Systema Naturae per Regna Tria Naturae, secundum Classes, Ordines,*
807 *Genera, Species, cum Characteribus, Differentiis, Synonymis, Locis. Laurenti Salvii,*
808 *Holmiae* 1, 1–823.

809 Longrich, N.R., Vinther, J., Pyron, R.A., Pisani, D., Gauthier, J.A., 2015. Biogeography of
810 worm lizards (Amphisbaenia) driven by end-Cretaceous mass extinction. *Proceedings*
811 *of the Royal Society B: Biological Sciences* 282, 20143034.

812 Martin, J.E., 2010. A new species of *Diplocynodon* (Crocodylia, Alligatoroidea) from the
813 Late Eocene of the Massif Central, France, and the evolution of the genus in the
814 climatic context of the Late Palaeogene. *Geological Magazine* 147, 596–610.

815 Martin, J.E., Gross, M., 2011. Taxonomic clarification of *Diplocynodon* Pomel, 1847
816 (Crocodylia) from the Miocene of Styria, Austria. *Neues Jahrbuch für Geologie und*
817 *Paläontologie–Abhandlungen* 261, 177–193.

818 Ménouret, B., Mein, P., 2008. Les vertébrés du miocène supérieur de Soblay (Ain, France)
819 Document des Laboratoires de Géologie de Lyon, UFR des sciences de la terre,
820 Université Claude Bernard-Lyon 1 165, 1–97.

821 Młynarski, M., 1976. Testudines, in: Kuhn, O., (Ed.), Encyclopedia of Paleoherpetology, part
822 7. G. Fischer Verlag, Stuttgart.

823 Oppel, M., 1811. Die Ordnungen, Familien, und Gattungen der Reptilien als Prodrom einer
824 Naturgeschichte derselben. Joseph Lindauer, München.

825 Pamouktchiev, A., de Lapparent de Broin, F., Anguelov, A., 1998. Nouvelle trouvaille de
826 tortue fossile en Bulgarie. Annuaire de l'Université de Sofia "St. Kliment Ohridski"
827 Faculté de Géologie et Géographie, Livre 1, Géologie 90, 5–8.

828 Paraskevaidis, I., 1955. Zwei Schildkröten reste aus dem Obermiozän von Chios. Annales
829 Géologiques des Pays Helléniques 6, 133–141.

830 Poulakakis, N., Lymberakis, P., Tsigenopoulos, C.S., Magoulas, A., Mylonas, M., 2005.
831 Phylogenetic relationships and evolutionary history of snake-eyed skink *Ablepharus*
832 *kitaibelii* (Sauria: Scincidae). Molecular Phylogenetics and Evolution 34, 245–256.

833 Purschke, C.A., 1885. *Clemmys sarmatica* n. sp. aus dem Tegel von Hernals bei
834 Wien. Denkschriften der Kaiserlichen Akademie der Wissenschaften / Mathematisch-
835 Naturwissenschaftliche Classe 50, 185–192.

836 Pyron, R.A., Reynolds, R.G., Burbrink, F.T., 2014. A taxonomic revision of boas (Serpentes,
837 Boidae). Zootaxa 3846, 249–260.

838 Pyron, R.A., Wiens, J.J., 2011. A large-scale phylogeny of Amphibia including over 2,800
839 species, and a revised classification of extant frogs, salamanders, and caecilians.
840 Molecular Phylogenetics and Evolution 61, 543–583.

841 Rage J.-C., 1984. Serpentes, in: Wellnhofer, P., (Ed.), Encyclopedia of Paleoherpetology, part
842 11. G. Fischer Verlag, Stuttgart, New York.

843 Rage, J.-C., 2003. Squamate reptiles from the early Miocene of Arrisdrift (Namibia). Memoir
844 of the Geological Society of Namibia (Ministry of Mines and Energy), Windhoek 19,
845 43–50.

846 Rage, J.-C., 2013. Mesozoic and Cenozoic squamates of Europe. *Palaeobiodiversity and*
847 *Palaeoenvironments* 93, 517–534.

848 Rage, J.-C., Augé, M., 1993. Squamates from the Cenozoic of the western part of Europe. A
849 review. *Revue de Paléobiologie*, volume spécial 7, 199–216. Rage, J.-C., Roček, Z.,
850 2003. Evolution of anuran assemblages in the Tertiary and Quaternary of Europe, in
851 the context of palaeoclimate and palaeogeography. *Amphibia-Reptilia* 24, 133–177.

852 Rage, J.-C., Sen, S., 1976. Les Amphibiens et les Reptiles du Pliocène supérieur de Calta
853 (Turquie). *Géologie méditerranéenne* 3, 127–134.

854 Richter, A., 1995. The vertebrate locality Maramena (Macedonia, Greece) at the Turolian–
855 Ruscinian Boundary (Neogene). 3. Lacertilia (Squamata, Reptilia). *Münchner*
856 *Geowissenschaftliche Abhandlungen* 28, 35–38.

857 Roček, Z., 1984. Lizards (Reptilia, Sauria) from the lower Miocene locality Dolnice
858 (Bohemia, Czechoslovakia). *Rozpravy Československé Akademie Ved, Rada*
859 *Matematických a přírodních Ved* 94, 3–69.

860 Rögl, F., 1999. Mediterranean and Paratethys. Facts and hypotheses of an Oligocene to
861 Miocene paleogeography (short overview). *Geologia Carpathica* 50, 330–349.

862 Römer, F., 1870. Über *Python euboicus*, eine fossile Riesenschlange aus tertiärem
863 Kalkschiefer von Kumi auf der Insel Euboea. *Zeitschrift der Deutschen Geologischen*
864 *Gesellschaft* 22, 582–590.

865 Rook, L., Delfino, M., Sami, M., 2015. I vertebrati fossili della cava del Monticino di
866 Brisighella: una finestra sui popolamenti continentali del Mediterraneo nel Miocene

867 superiore. I gessi di Brisighella e Rontana. Memorie dell'Istituto Italiano di Speleologia
868 s. II 28, 79–100.

869 Rustioni, M., Mazza, P., Azzaroli, A., Boscagli, G., Cozzini, F., Di Vito, E., Masseti, M.,
870 Pisano, M., 1993. Miocene vertebrate remains from Scontrone, National Park of
871 Abruzzi, Central Italy. Rendiconti della Accademia Nazionale dei Lincei, Classe di
872 Scienze Fisiche Matematiche e Naturali 9, 227–237.

873 Sanchiz, B., 1998. Salientia, in: Wellnhofer, P., (Ed.), Encyclopedia of Paleoherpetology, part
874 4. Friedrich Pfeil, Munich.

875 Sen, S., Antoine, P.O., Varol, B., Ayyildiz, T., Sözeri, K., 2011. Giant rhinoceros
876 *Paraceratherium* and other vertebrates from Oligocene and middle Miocene deposits of
877 the Kağızman-Tuzluca Basin, Eastern Turkey. Naturwissenschaften 98, 407–423.

878 Sindaco, R., Jeremčenko, V., 2008. The reptiles of the western Palearctic. Vol. 1: annotated
879 checklist and distributional atlas of the turtles, crocodiles, amphisbaenians and lizards of
880 Europe, North Africa, Middle East and Central Asia. Monografie della Societas
881 Herpetologica Italica I, Belvedere, Latina.

882 Sindaco, R., Kornilios, P., Sacchi, R., Lymberakis, P., 2014: Taxonomic reassessment of
883 *Blanus strauchi* (Bedriaga, 1884) (Squamata: Amphisbaenia: Blanidae), with the
884 description of a new species from south-east Anatolia (Turkey). Zootaxa 3795, 311–
885 326.

886 Staesche, K., Karl, H.-V., Staesche, U., 2007. Fossile Schildkröten aus der
887 Türkei. Geologisches Jahrbuch Reihe B 98, 91–149.

888 Steel, R., 1973. Crocodylia, in: Kuhn, O., (Ed.), Encyclopedia of Paleoherpetology, part 16.
889 G. Fischer Verlag, Stuttgart, Portland.

890 Szalai, T., 1931. Schildkrötenstudien: I. *Testudo schafferi* nov. sp., eine Riesenschildkröte aus
891 dem Pliozän von Samos. Annalen des Naturhistorischen Museums in Wien 46, 153–157.

892 Szyndlar, Z., 1991a. A review of Neogene and Quaternary snakes of Central and Eastern
893 Europe. Part I: Scolecophidia, Boidae, Colubrinae. *Estudios Geológicos* 47, 103–126.

894 Szyndlar, Z., 1991b. A review of Neogene and Quaternary snakes of Central and Eastern
895 Europe. Part II: Natricinae, Elapidae, Viperidae. *Estudios Geológicos* 47, 237–266.

896 Szyndlar, Z., 1995. The vertebrate locality Maramena (Macedonia, Greece) at the Turolian–
897 Ruscinian Boundary (Neogene). 4. Serpentes (Squamata, Reptilia). *Münchner*
898 *Geowissenschaftliche Abhandlungen* 28, 35–39.

899 Szyndlar, Z., Rage, J.-C., 2002. Fossil record of the true vipers, in: Schuett, G.W., Höggren,
900 M., Douglas, M.E., Greene, H.W., (Eds.), *Biology of the vipers*. Eagle Mountain
901 Publishing, Eagle Mountain, pp. 419–444.

902 Szyndlar, Z., Rage, J.-C., 2003. Non-erycine Booidea from the Oligocene and Miocene of
903 Europe. Institute of Systematics and Evolution of Animals, Polish Academy of Sciences,
904 Cracow.

905 Szyndlar, Z., Zerova, G., 1990. Neogene cobras of the genus *Naja* (Serpentes: Elapidae) of
906 East Europe. *Annalen des Naturhistorischen Museums in Wien* 91A, 53–61.

907 Theobald, W., Jr., 1868. Catalogue of Reptiles in the Museum of the Asiatic Society of
908 Bengal. *Journal of the Asiatic Society*, Extra Number.

909 Valakos, E.D., Pafilis, P., Sotiropoulos, K., Lymberakis, P., Maragou, P., Foufopoulos, J.,
910 2008. *Amphibians and Reptiles of Greece*. Edition Chimaira, Serpents Tale, Frankfurt.

911 Valenciennes, A., 1833. *Chelonia pelasgorum*, *Emys hellenica*, *Emys iberica*, *Emys rivulata*,
912 in: Bory de Saint-Vincent, J.B., (Ed.), *Expédition Scientifique de Morée*. Travaux de la
913 Section des Sciences Physiques. Zoologie. Levrault, F.G., Paris.

914 van der Made, J., 1996. Pre-Pleistocene land mammals from Crete, in: Reese, D.S., (Ed.),
915 *Pleistocene and Holocene fauna of Crete and its first settlers*. Monographs in World
916 Archaeology 28, Prehistory Press, pp. 69–79.

917 Venczel, M., Știucă, E., 2008. Late middle Miocene amphibians and squamate reptiles from
 918 Taut, Romania. *Geodiversitas* 30, 731–763.

919 Vianna, A., Moraes, A., 1945. Sur un craîne de crocodile fossile découvert dans le
 920 Miocène de Lisbonne. *Boletim de la Sociedade Geológica de Portugal* 4, 161–170.

921 Viriot, L., Pelaez-Campomanes, P., Vignaud, P., Andossa, L., Mackaye, H.T., Brunet, M.,
 922 2011. A new xerinae (Rodentia, Sciuridae) from the late Miocene of Toros-Menalla
 923 (Chad). *Journal of Vertebrate Paleontology* 31, 844–848.

924 Vitek, N.S., Joyce, W.G., 2015. A review of the fossil record of New World turtles of the
 925 clade *Pan-Trionychidae*. *Bulletin of the Peabody Museum of Natural History* 56, 185–
 926 244.

927 Vitt, L.J., Caldwell, J.P., 2014. *Herpetology. An introductory biology of amphibians and*
 928 *reptiles. Fourth edition. Academic Press, Elsevier, Amsterdam, Boston, Heidelberg,*
 929 *London, New York, Oxford, Paris, San Diego, San Francisco, Sydney, Tokyo.*

930 Vlachos, E., 2015. The fossil chelonians of Greece. Systematics - evolution - stratigraphy -
 931 palaeoecology. Ph.D. thesis, Aristotle University of Thessaloniki, Greece.

932 Vlachos, E., Cerda, I., Tsoukala, E., 2015a. The first record of a soft-shelled turtle
 933 (Testudines: *Pan-Trionychidae*) from southern Balkans (Pliocene, Gefira, N. Greece)
 934 and new information from bone histology. *The Science of Nature* 102, 45.

935 Vlachos, E., Kotsakis, T., Delfino, M., 2015b. The chelonians from the latest Miocene–earliest
 936 Pliocene localities of Allatini and Pylea (East Thessaloniki, Macedonia, Greece).
 937 *Comptes Rendus Palevol* 14, 187–205.

938 Vlachos, E., Tsoukala, E., 2014. *Testudo* cf. *graeca* from the new Late Miocene locality of
 939 Platania (Drama basin, N. Greece) and a reappraisal of previously published specimens.
 940 *Bulletin of the Geological Society of Greece* XLVIII, 27–40.

Wallach, V., Williams, K.L., Boundy, J., 2014. Snakes of the world: a catalogue of living and extinct species. CRC Press, Boca Raton.

Weithofer, A., 1888. Beiträge zur Kenntniss der Fauna von Pikermi bei Athen. Beiträge zur Paläontologie Österreich-Ungarns 6, 225–292.

Werner, F., 1898. Über einige neue Reptilien und einen neuen Frosch aus dem cilicischen Taurus. Zoologischer Anzeiger 21, 217–223.

Zangerl, R., 1969. The turtle shell, in: Bellairs d' A.A., Parsons, T.S., (Eds.), The Biology of the Reptilia 1, Morphology A. Academic Press, London, New York, pp. 311–339.

Figure captions




Fig. 1. Map of Crete, indicating the earliest late Miocene locality of Plakias.



Fig. 2. cf. Alytidae from the earliest late Miocene of Plakias. UU PL 701, a trunk vertebra in dorsal (A), anterior (B) and ventral(C) views.




Fig. 3. Pan-Trionychidae from the earliest late Miocene of Plakias. A, NHMC 21.7.3.1670, fragment of a neural, in A1, dorsal and A2, visceral views. B, UU PL 703, fragment of a costal in B1, dorsal and B2, visceral views. C, UU PL 704, fragment of a costal in C1, dorsal and C2, visceral views. D, NHMC 21.7.3.1671, fragment of plastron in dorsal view.




Fig. 4. *Mauremys* sp. from the earliest late Miocene of Plakias. A, NHMC 21.7.3.1623, neural in (A1) dorsal and (A2) visceral views. B, NHMC 21.7.3.1622, neural in (B1) dorsal, (B2) drawing of the dorsal and (B3) visceral views. C, NHMC 21.7.3.1645, neural in (C1) dorsal and (C2) visceral views. D, NHMC 21.7.3.1624, neural in (D1) dorsal and (D2) visceral

969 views.E, NHMC 21.7.3.1651, neural in (E1) dorsal, (E2) drawing of the dorsal and (E3)
970 visceral views. F, NHMC 21.7.3.1649, neural fragment in (F1) dorsal (F3) visceral views. G,
971 NHMC 21.7.3.1640, two successive costals in (G1) dorsal and (G2) visceral views.H, NHMC
972 21.7.3.1638, costal in (H1) dorsal, (H2) drawing of the dorsal and (H3) visceral views. I,
973 NHMC 21.7.3.1639, costal fragment in (I1) dorsal, (I2) drawing of the dorsal and (I3) visceral
974 views. J, NHMC 21.7.3.1641, peripheral in (J1) dorsal and (J2) drawing of the dorsal views.
975 K, NHMC 21.7.3.1618, right epiplastron and entoplastron in (K1) visceral, (K2) ventral and
976 (K3) drawing of the ventral views. L, NHMC 21.7.3.1637, left epiplastron in (L1) visceral,
977 (L2) ventral and (L3) drawing of the ventral views. M, NHMC 21.7.3.1635, right epiplastron
978 in (M1) visceral, (M2) ventral and (M3) drawing of the ventral views. N, NHMC 21.7.3.1632,
979 left epiplastron fragment in (N1) visceral, (N2) ventral and (N3) drawing of the ventral views.
980 O, NHMC 21.7.3.1620, left hypoplastron in (O1) visceral, (O2) ventral and (O3) drawing of
981 the ventral views. P, NHMC 21.7.3.1643, right hypoplastron in (P1) visceral, (P2) ventral and
982 (P3) drawing of the ventral views. Q, NHMC 21.7.3.1619, right xiphiplastron in (Q1)
983 visceral, (Q2) ventral and (Q3) drawing of the ventral views. R, NHMC 21.7.3.1642 right
984 xiphiplastron fragment in (R1) visceral and (R2) ventral views. S, UU PL 705, possible
985 hyoplastron fragment in (S1) visceral, (S2) ventral and (S3) drawing of the ventral views.
986 Abbreviation: ir, irregular scute growth.

987




Fig. 5. Crocodylia from the earliest late Miocene of Plakias. A, UU PL 706, isolated tooth in labial (A1) and mesial (A2) views. B, UU PL 707, isolated tooth in labial (B1) and mesial (B2) views. C, UU PL 708, isolated tooth in labial (C1) and mesial (C2) views. D, UU PL 709, isolated tooth in labial (D1) and mesial (D2) views.




Fig. 6. Amphisbaenia from the earliest late Miocene of Plakias. UU PL 733, presacral vertebra in right lateral (A), dorsal (B), ventral (C) and anterior (D) views.




Fig. 7. ?Natricinae from the earliest late Miocene of Plakias. UU PL 734, preloacal vertebra in dorsal (A), anterior (B), right lateral (C), posterior (D) and ventral (E) views.

Table 1

The youngest occurrences of crocodylians in Europe.

Taxon	Age	Locality	Reference
<i>Crocodylus</i> sp.	latest Messinian (or even early Zanclean,	Gargano, Apulia, Italy	Delfino et al., 2007

	depending on the age		
	of the Terre Rosse		
	from Gargano)		
Crocodylia indet.	late Messinian	Cava del Monticino, Brisighella, Emilia- Romagna, Italy	Rook et al., 2015
<i>Tomistoma</i> <i>calaritanus</i>	Tortonian–Messinian	Is Mirrionis, Sardinia, Italy	Capellini, 1890
Crocodylia indet.	late Tortonian	Fiume Santo, Sardinia, Italy	Abbazzi et al., 2008
cf. <i>Crocodylus</i> sp. (type of <i>Crocodylus</i> <i>bambolii</i>)	Tortonian	Montabamboli- Casteani-Ribolla, Tuscany, Italy	Delfino and Rook, 2008
<i>Tomistoma</i> cf. <i>lusitanica</i>	Tortonian	Olhos de Agua, Algarve, Portugal	Vianna and Moraes, 1945
Crocodylia indet. (<i>Diplocynodon</i> sp.)	Tortonian	Soblay, Ain, France	Ménouret and Mein, 2008
Crocodylia indet.	early Tortonian	Plakias, Crete, Greece	this paper
cf. <i>Crocodylus</i> sp.	early Tortonian	Scontrone, Abruzzo, Italy	Rustioni et al., 1993; Delfino and Rossi, 2013

